

References on QFT for critical phenomena and Wilsonian ideas

- Chapters 12 and 13 of Peskin's book.
- Cardy's book "Scaling and renormalization in statistical physics".
- David Tong's lectures on Statistical Field Theory, <http://www.damtp.cam.ac.uk/user/tong/sft.html>
- Silviu Pufu's lectures at the São Paulo bootstrap school, <http://bootstrap.ictp-saifr.org/school/>
- David Simmons-Duffin's lectures (chapters 1-3) <https://github.com/davidsd/ph229/blob/master/ph229-notes.pdf>
- Wilson and Kogut's Physics Reports, pdf linked from the webpage. I highly recommend reading Section 12, on the topology of the RG and the *definition* of QFT.
- Polchinski's paper "Renormalization and effective Lagrangians", pdf linked from the webpage. The introduction and the toy model of section 2 are accessible and highly recommended reading. It should be clear that the inspiration for this paper came from Section 12 of Wilson and Kogut. Weinberg gives a simplified summary of Polchinski's argument in section 12.4 of volume 1 of his QFT textbook.

Critical exponents versus conformal dimensions cheat sheet

We are going to relate the critical exponents α , β , γ , δ , η and ν to the conformal dimensions Δ_ϕ and Δ_{ϕ^2} . This analysis applies to systems with a two-dimensional phase diagram, *i.e.*, with two relevant scalar operators, ϕ and ϕ^2 . We use the terminology appropriate to magnetic systems: ϕ couples to the magnetic field h and ϕ^2 couples to $m^2 \sim t$, where t is the reduced temperature, $t = (T - T_c)/T_c$. We indicate by f the free energy per unit volume, which has mass dimension d .

- Exponent η :

Two-point function of the fundamental scalar

$$\langle \phi(x)\phi(0) \rangle = \frac{1}{|x|^{2\Delta_\phi}} \quad \Delta_\phi = \frac{d}{2} - 1 + \gamma_\phi,$$

where γ_ϕ is the “anomalous dimension” of ϕ , *i.e.* the deviation from the naive engineering dimension. Comparing with

$$\langle \phi(x)\phi(0) \rangle = \frac{1}{|x|^{d-2+\eta}}$$

we find

$$\eta = 2\gamma_\phi.$$

- Exponent ν :

Defined from the behavior of the correlation length,

$$\xi \sim \frac{1}{|T - T_c|^\nu}.$$

Dimensional analysis gives

$$\xi \sim \frac{1}{(m^2)^{\frac{1}{[m^2]}}} \sim \frac{1}{t^{\frac{1}{[m^2]}}}.$$

The composite operator ϕ^2 has dimension $\Delta_{\phi^2} = 2(\frac{d}{2} - 1) + \gamma_{\phi^2}$. Since ϕ^2 appears as $m^2\phi^2$ in the action, we have

$$[m^2] + \Delta_{\phi^2} = d$$

which implies

$$[m^2] = 2 - \gamma_{\phi^2}, \quad [m] = 1 - \frac{1}{2}\gamma_{\phi^2}.$$

This gives

$$\nu = \frac{1}{[m^2]} = \frac{1}{2 - \gamma_{\phi^2}}.$$

Recall also that $\gamma_{\phi^2} = 2\gamma_m$ (see solutions of homework 1).

- Exponent α :

$$\frac{\partial^2 f}{\partial t^2} \sim |t|^{-\alpha}$$

Dimensional analysis gives

$$\alpha = -\frac{d - 2[m^2]}{[m^2]} = 2 - \frac{d}{2 - \gamma_{\phi^2}}$$

- Exponent β :

$$\lim_{h \rightarrow 0^+} \langle \phi \rangle \sim (-t)^\beta.$$

Dimensional analysis gives

$$\beta = \frac{\Delta_\phi}{[m^2]} = \frac{d/2 - 1 + \gamma_\phi}{2 - \gamma_{\phi^2}}.$$

- Exponent γ :

$$\frac{\partial^2 f}{\partial h^2} \sim |t|^{-\gamma}$$

Dimensional analysis gives

$$\gamma = -\frac{d - 2[h]}{[m^2]} = \frac{2(d - \Delta_\phi) - d}{2 - \gamma_\phi^2} = \frac{2 - 2\gamma_\phi}{2 - \gamma_\phi^2}$$

- Exponent δ :

$$\langle \phi \rangle \sim h^{1/\delta}$$

Dimensional analysis gives

$$\delta = \frac{[h]}{\Delta_\phi} = \frac{d/2 + 1 - \gamma_\phi}{d/2 - 1 + \gamma_\phi}.$$

The “scaling relations”

$$\gamma = \nu(2 - \eta), \quad \alpha + 2\beta + \gamma = 2, \quad \delta = 1 + \frac{\gamma}{\beta}, \quad \alpha = 2 - \nu d$$

are automatically obeyed.